

# SPECT*(i)*CLE™ G

Preemergent Herbicide for the Control of Annual Grasses, Annual Sedges, and Broadleaf Weeds in Turfgrass, Landscape Ornamentals, and Hard-scapes

**DO NOT USE FOR THE MANUFACTURING OF FERTILIZER**

**ACTIVE INGREDIENT:**

INDAZIFLAM .....0.0224%

**OTHER INGREDIENTS:** .....99.9776%

Total: .....100.000%

EPA Reg. No. 432-1523

Contains 0.0112 lbs of INDAZIFLAM in a 50 lb bag

## KEEP OUT OF REACH OF CHILDREN

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A  
Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-331-2867

### FIRST AID

|                                                                                                                     |                                                                                                                                                                                                                                                                                                                                                     |
|---------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>If swallowed:</b>                                                                                                | <ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul> |
| <b>If in eyes:</b>                                                                                                  | <ul style="list-style-type: none"> <li>• Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>                                                           |
| <b>If on skin or clothing:</b>                                                                                      | <ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>                                                                                                               |
| <b>If inhaled:</b>                                                                                                  | <ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>                                             |
| Have the product container or label with you when calling a poison control center or doctor or going for treatment. |                                                                                                                                                                                                                                                                                                                                                     |

See inside leaflet for complete First Aid Instructions, Precautionary Statements, Directions for Use and Storage and Disposal Instructions.

Net Weight:

**50 lbs**

**80849265**

80946255A

121019AV3

**BACKED**  
by **BAYER™**

## PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## ENGINEERING CONTROLS STATEMENTS:

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing and PPE immediately if pesticide gets inside. Then wash body thoroughly and put on clean clothing.

## ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean watermark. Do not contaminate water when disposing of equipment rinseate or washwater. This product may enter water through runoff. Follow directions for use to avoid runoff. A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for getting into water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

**Surface Water Advisory:** This pesticide may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

**Ground Water Advisory:** This pesticide has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

## DIRECTIONS FOR USE

**It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**Read entire label before using this product.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, waterproof gloves and shoes plus socks

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Always wear protective clothing and use appropriate equipment during application and handling. Avoid contact with granules. Avoid breathing dust.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

### PESTICIDE STORAGE

Store in original container only. Store in a manner to prevent cross-contamination of other pesticides, fertilizer, food, and feed. Do not store in or around the home. Store in a dry, locked storage area.

### PESTICIDE DISPOSAL

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

### CONTAINER HANDLING

Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Offer for recycling, if available. Dispose of in a sanitary landfill, or by incineration; or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## PRODUCT INFORMATION

**SPECTICLE G** is a selective, preemergent alkylazine herbicide. **SPECTICLE G** provides extended residual control of many annual grasses including crabgrass, goosegrass, and annual bluegrass, as well as annual sedges, and some broadleaf weeds in turf, landscape ornamentals, non-bearing fruit and nut trees in residential plantings, as well as hardscapes.

Use **SPECTICLE G** on established turf in areas including golf courses (roughs and fairways), residential and commercial lawns, parks, and cemeteries.

**SPECTICLE G** controls weeds by reducing the emergence of seedlings through inhibition of cellulose biosynthesis (CB Inhibitor). Necrosis or yellowing may be observed if the herbicide is applied to herbaceous tissue such as leaves and green stems of susceptible plants including sensitive ornamentals. **SPECTICLE G** does not control plants emerging from tubers, rhizomes, bulbs,

corms, existing rootstocks, and woody vegetation. **SPECTICLE G** suppresses the development of some additional hard-to-control weeds.

**SPECTICLE G** needs to be activated prior to weed germination for most effective control. For maximum activity against germinating weeds, **SPECTICLE G** requires rainfall or irrigation prior to weed germination.

**SPECTICLE G** is a very active herbicide that provides effective weed control when applied to the soil around many woody oramentals grown in-ground. **SPECTICLE G** may damage sensitive plants, if granules are allowed to remain in contact with foliage. Use **SPECTICLE G** according to the label.

| SPECTICLE G Conversion Table |                        |
|------------------------------|------------------------|
| Rate in lbs/Acre             | Rate in lbs/1000 sq ft |
| 100                          | 2.3                    |
| 125                          | 2.9                    |
| 150                          | 3.4                    |
| 200                          | 4.6                    |
| 400                          | 9.2                    |

PRODUCT USE RESTRICTIONS

- Do not exceed 200 lbs of **SPECTICLE G** lbs per acre (4.6 lbs/1000 sq ft) in a single application for turf and ornamental applications.
- Do not exceed 400 lbs of **SPECTICLE G** lbs per acre (9.2 lbs/1000 sq ft) for all applications within a 12-month period.
- Do not contaminate water intended for irrigation and domestic use.
- Do not apply using aerial application equipment.

DEACTIVATING SPECTICLE G

Activated charcoal has been shown to deactivate **SPECTICLE G** if applied within several days of an application. Application of activated charcoal within 2 weeks after an application of **SPECTICLE G** will not reverse phytotoxic symptoms immediately, but it will aid in recovery over time. If it should be necessary to re-sod areas treated with **SPECTICLE G**, remove damaged turf to a depth of at least 2 inches, cultivate the soil, and apply activated charcoal to bare ground prior to laying sod. Follow directions for the amount of charcoal to apply on the label of the activated charcoal.

APPLICATION INFORMATION

Apply **SPECTICLE G** with clean, properly calibrated drop, rotary, or hand shaker equipment according to the manufacturer's directions. Check periodically to be certain that the equipment is working properly prior to each use. Uniform application is essential for satisfactory weed control. Follow instructions in the table below for proper calibration of equipment. Avoid skips for the best weed control and overlapping application patterns to avoid turf injury.

CALIBRATION SETTINGS FOR COMMONLY USED SPREADERS

HOW TO CALIBRATE SPREADERS FOR APPLICATION OF SPECTICLE G

The spreader settings provided are accurate for the product samples tested. Results obtained by users may vary due to spreader model variables, spreader calibration differences, granular product production, operator-induced variables such as walking speed and handle height, and also by environmental factors such as air temperature and relative humidity. The settings provided below are recommended as a starting point for accurate calibration.

| SPECTICLE G Rates                                  |                                    |                                    |
|----------------------------------------------------|------------------------------------|------------------------------------|
| Rotary Spreaders                                   | 150 lbs/acre<br>3.4 lbs/1000 sq ft | 200 lbs/acre<br>4.6 lbs/1000 sq ft |
| Scotts Rotary                                      | I                                  | J                                  |
| Lesco Rotary                                       | C.5                                | D.5                                |
| Spyker 288                                         | 3                                  | 3.5                                |
| Drop Spreaders                                     | 150 lbs/acre<br>3.4 lbs/1000 sq ft | 200 lbs/acre<br>4.6 lbs/1000 sq ft |
| Lesco Drop                                         | 3.25                               | 3.5                                |
| Handheld Spreaders *                               | 150 lbs/acre<br>3.4 lbs/1000 sq ft | 200 lbs/acre<br>4.6 lbs/1000 sq ft |
| * Calibration run in traditional calibration trays |                                    |                                    |
| Solo 421 S                                         | 3.25                               | 4.5                                |
| Earthway Ev-N                                      | 10.5                               | 12                                 |
| Warren T-7                                         | 5.5                                | 6                                  |
| Spyker                                             | 3                                  | 3.5                                |
| Handheld Spreaders **                              | 150 lbs/acre<br>3.4 lbs/1000 sq ft | 200 lbs/acre<br>4.6 lbs/1000 sq ft |
| ** Calibration run in round ornamental containers  |                                    |                                    |
| Solo 421 S                                         | 3                                  | 3.5                                |
| Earthway Ev-N                                      | 8                                  | 9                                  |
| Warren T-7                                         | 3.5                                | 4                                  |
| Spyker                                             | 2.75                               | 3.25                               |

## SPECTICLE G Rates (continued)

| Tractor Mount Spreaders | 150 lbs/acre<br>3.4 lbs/1000 sq ft | 200 lbs/acre<br>4.6 lbs/1000 sq ft |
|-------------------------|------------------------------------|------------------------------------|
| Vicon                   | 16                                 | 20                                 |
| Lely                    | 4.75                               | 5.5                                |

### IRRIGATION AND WATERING AFTER APPLICATION

To activate **SPECTICLE G** for maximum herbicidal benefit, irrigate within 2 days after application to turf, ornamentals, or other uses to move the herbicide into the soil. With dry soil, use 0.25 inch of irrigation water, and with soil at or greater than field capacity, reduce the amount of irrigation water. In either case, do not create conditions that cause visible run-off of irrigation water. Adequate rainfall following an application will negate the need for irrigation. For application to ornamentals over-the-top, irrigate within 1 hour to wash granules from the leaf surface and activate the product in the soil.

### EQUIPMENT CLEANUP PROCEDURE

Equipment used to apply **SPECTICLE G** must be cleaned prior to use on other sensitive turf and ornamentals, or injury may result. Before and after using **SPECTICLE G**, clean all equipment on an approved rinse pad. Wear proper PPE while cleaning equipment.

### RESISTANCE MANAGEMENT

**SPECTICLE G** controls weeds by inhibiting cellulose biosynthesis. **SPECTICLE G** is classified as a Group 29 herbicide. No known resistance to **SPECTICLE G** exists, and there are no known instances of cross-resistance between **SPECTICLE G** and other classes of herbicides with other modes of action. Performance of **SPECTICLE G** is not affected by the presence of biotypes resistant to glyphosate, triazines, ALS inhibitors, mitotic inhibitors, growth regulators, or herbicides with other modes of action. Prolonged use of any herbicide with the same mode of action may lead to selection of resistant weed populations and a reduction in efficacy. Rotation of herbicides with an alternate mode of action can help prevent the development and spread of resistant weed populations. If resistance to **SPECTICLE G** is suspected, use a herbicide with an alternate mode of action. The decision to use any particular herbicide should be based on an IPM program and all cultural practices and other pesticides in use.

## DIRECTIONS FOR USE OF SPECTICLE G ON TURF

### USE RESTRICTIONS FOR SPECTICLE G ON TURF

- Do not apply **SPECTICLE G** to newly seeded turf. Established turf is defined as turf at least 16 months after seeding or sprigging
- Do not apply to golf course greens, tees, or collars.
- Do not verticut within 1 month before or after an application of **SPECTICLE G**.
- Do not apply to turf that is showing significant signs of stress including winterkill, spring dead spot, transitioning ryegrass, Bermudagrass spring root decline, soil compaction, foliar and root diseases, nematodes, salt accumulation, shade, excessive foot or equipment traffic, and drought. Application of **SPECTICLE G** in these situations may delay turf recovery.

### USE OF SPECTICLE G ON WEAKENED OR STRESSED TURF

Do not apply **SPECTICLE G** to weakened turf that requires significant fill-in or recovery. **SPECTICLE G** controls weeds by inhibiting root development. Newly developing roots of desirable turf may be affected. Factors, both biotic and abiotic, that weaken turf include winterkill, spring dead spot, transitioning ryegrass, Bermudagrass spring root decline, soil compaction, foliar and root diseases, nematodes, salt accumulation, shade, excessive foot traffic, and drought. Application of **SPECTICLE G** to turf in these situations may delay turf recovery. **SPECTICLE G** may also be used on dormant turf. The user should assess the condition of turf prior to dormancy and determine whether an application of **SPECTICLE G** is appropriate.

Application of **SPECTICLE G** to stressed turf may produce symptoms of injury including yellowing, purple discoloration, thinning, and necrosis. If injury occurs, promote recovery through optimizing fertility and other cultural practices.

### TURF TOLERANCE

**SPECTICLE G** can be used on the following turf types.

#### Warm season turfgrasses and their cultivars

- **Bermudagrass\*** (*Cynodon dactylon*) (and hybrids) - *Baby, Celebration, Common\*\**, *Floralex, GN1, TifSport, Tifton 10, Tifway 419, Princess, Vamont, Riviera, Yukon*
  - **Zoysiagrass** (*Zoysia* spp.) - *Cavalier, Crowne, DeAnza, El Toro, Empire, GNZ, Jamar, Meyer, Palisades, Zenith, Zeon*
  - **Centipedegrass\*** (*Eremochloa ophiuroides*) - *Tifblair*
  - **St. Augustinegrass\*** (*Stenotaphrum secundatum*) - *Captiva, Floratam, Floratine, Palmetto, Raleigh*
  - **Buffalograss** (*Buchloe dactyloides*)
  - **Bahiagrass** (*Paspalum notatum*)
- \* Application of **SPECTICLE G** may temporarily inhibit the rooting of stolons of bermudagrass, centipedegrass, and St. Augustinegrass.

\*\* Common or off-type Bermudagrass may show increased sensitivity to **SPECTICLE G** compared to hybrids. Use lower application rates as indicated in the table below.

Before making a large-scale application to other cultivars, treat a small area of turf with **SPECTICLE G** and allow four weeks to assess acceptable turf tolerance.

Maximum Single Application Rates for SPECTICLE G on Warm Season Grasses

| Turf Type                   | SPECTICLE G Rate<br>(lbs/acre) | SPECTICLE G Rate<br>(lbs/1000 sq feet) |
|-----------------------------|--------------------------------|----------------------------------------|
| Bermudagrass (Hybrids)      | 200                            | 4.6                                    |
| Bermudagrass (Common types) | 125                            | 2.9                                    |
| Zoysiagrass                 | 200                            | 4.6                                    |
| Buffalograss                | 200                            | 4.6                                    |
| Bahiagrass                  | 200                            | 4.6                                    |
| St. Augustinegrass          | 100                            | 2.3                                    |
| Centipedegrass              | 100                            | 2.3                                    |

SPECTICLE G may cause unacceptable injury to some turf types. **Do not use** on the following turf types or on turf that is a mixture containing any of these grasses, unless thinning or removal is desired.

- **Creeping bentgrass** (*Agrostis palustris*)
  - **Colonial bentgrass** (*Agrostis capillaris*)
  - **Annual bluegrass** (*Poa annua*)
  - **Roughstalk bluegrass** (*Poa trivialis*)
  - **Fine fescue** (*Festuca rubra*)
  - **Turf type tall fescue** (*Festuca arundinaceae*)
- **Kikuyugrass** (*Pennisetum clandestinum*)
  - **Perennial ryegrass** (*Lolium perenne*)
  - **Annual ryegrass** (*Lolium multiflorum*)
  - **Kentucky bluegrass** (*Poa pratensis*)
  - **Seashore paspalum** (*Paspalum vaginatum*)
  - **Bermudagrass** (*Cynodon dactylon*) Common variety "Ormond"

PRECAUTIONS FOR USE OF SPECTICLE G NEAR SENSITIVE GRASSES

**CAUTION: SPECTICLE G can cause turf injury and stand reduction to sensitive grasses listed in the Turf Tolerance section of this label.**

- **SPECTICLE G** may affect sensitive grasses downslope from treated areas after excessive rainfall. To minimize off-target effects of **SPECTICLE G** on sensitive grasses, irrigate after application as directed in the **IRRIGATION AND WATERING AFTER APPLICATION** section of this label and follow the directions below for specific situations.
- Allow turf to dry before allowing foot traffic or equipment through treated areas near sensitive grasses to avoid tracking.

PRECAUTIONS AROUND GREENS CONTAINING COOL SEASON GRASSES

- For applications of **SPECTICLE G** adjacent to bentgrass greens and Bermudagrass greens overseeded with roughstalk bluegrass (*Poa trivialis*) or other cool season grasses, follow the precautions above and provide an untreated buffer zone of at least **15 ft**.
- Do not apply **SPECTICLE G** on uphill slopes adjacent to bentgrass or Bermudagrass greens overseeded with cool season grasses such as *Poa trivialis*.

PRECAUTIONS FOR BERMUDAGRASS OVERSEEDED WITH RYEGRASS

- **Do not apply SPECTICLE G** on annual and perennial ryegrass or on Bermudagrass overseeded with ryegrass. Provide an untreated buffer zone of at least **15 ft** between overseeded Bermudagrass and treated areas.

PRECAUTIONS FOR USE NEXT TO OTHER SENSITIVE GRASSES

- **SPECTICLE G** may be applied in situations where warm season grasses are adjacent to sensitive grasses such as tall fescue and perennial ryegrass. The applicator, however, must take care not to apply **SPECTICLE G** directly to sensitive grasses or injury may occur. Cool season grasses must be established at least 16 months prior to an application of **SPECTICLE G** on an adjacent warm season grass. Maintain an untreated buffer of **15 feet** between sensitive grasses and treated areas.

USE RATES, TIMINGS, AND MAXIMUM SEASONAL RATE FOR SPECTICLE G ON TURF

The desired rate of **SPECTICLE G** for each application and the interval between applications depends on the turfgrass being treated, the weeds to be controlled, and local conditions such as soil texture. Applications of **SPECTICLE G** are made in any season prior to weed germination. Depending on local conditions and weed control desired, sequential applications of **SPECTICLE G** may be made in a calendar year.

The initial application of **SPECTICLE G** must be 100-200 lbs of product per acre (2.3-4.6 lbs/1000 sq ft). Use higher rates within the rate range if weed pressure is expected to be high or when longer residual control is desired. The optimum single rate for each weed at a particular use site may be affected by previous practices, including history of herbicide use, mowing height, turf density, and turf health.

Soil texture influences the best use rate for **SPECTICLE G**. Use lower rates in coarse or sandy soils. In fine soils or in soils containing high amounts of organic matter higher use rates of **SPECTICLE G** may be used. The table below provides guidelines for appropriate rates.

SINGLE APPLICATION PROGRAM FOR SPECTICLE G\*

| Target Weed               | SPECTICLE G Rate<br>(lbs/acre)** | SPECTICLE G Rate<br>(lbs/1000 sq ft) |
|---------------------------|----------------------------------|--------------------------------------|
| Crabgrass                 | 100-200                          | 2.3-4.6                              |
| Goosegrass                |                                  |                                      |
| Annual Bluegrass          |                                  |                                      |
| Broadleaf weeds           |                                  |                                      |
| Annual sedge and kyllinga | 200                              | 4.6                                  |

\*Use higher use rates within the rate range for each target weed when a site has historically higher weed pressure. Also, use higher rates on fine textured soils and lower rates on medium and coarse textured soils.

\*\*The maximum single application on turf is 200 lbs per acre (4.6 lbs/1000 sq ft). Do not exceed 400 lbs of **SPECTICLE G** per acre (9.2 lbs/1000 sq ft) in a 12-month period.

Split or multiple applications of **SPECTICLE G** can be made in a use season to extend the length of control of specific weeds, to control weeds germinating over an extended period, and to control other weeds emerging during the same or overlapping period. Depending on the weed pest (see table below), make follow-up applications within 30-90 days after the initial application. The total amount of **SPECTICLE G** applied in a calendar year must not exceed 400 lbs of product per acre (9.2 lbs/1000 sq ft).

**SPLIT APPLICATION PROGRAM FOR SPECTICLE G**  
**Use rate for Split Applications of SPECTICLE G**

| Target Weed               | Rate Range for SPECTICLE G (initial application) |                   | Rate Range for SPECTICLE G (follow up application within 45-90 days) |                   |
|---------------------------|--------------------------------------------------|-------------------|----------------------------------------------------------------------|-------------------|
|                           | In lbs/acre                                      | In lbs/1000 sq ft | In lbs/acre                                                          | In lbs/1000 sq ft |
| Crabgrass                 | 100-200                                          | 2.3-4.6           | 100-200                                                              | 2.3-4.6           |
| Goosegrass                |                                                  |                   |                                                                      |                   |
| Annual Bluegrass          |                                                  |                   |                                                                      |                   |
| Broadleaf weeds           | 100-200                                          | 2.3-4.6           | 100-200                                                              | 2.3-4.6           |
| Annual sedge and kyllinga | 200                                              | 4.6               | 125-200                                                              | 2.9-4.6           |

**MULTIPLE APPLICATION PROGRAM FOR SPECTICLE G**  
**Use Rates for Multiple Applications of SPECTICLE G**

| Target Weed | Rate Range for SPECTICLE G (initial application) |     | Rate Range for SPECTICLE G (follow up application within 30-45 days) | Rate Range for SPECTICLE G (follow up application within 30-45 days) |
|-------------|--------------------------------------------------|-----|----------------------------------------------------------------------|----------------------------------------------------------------------|
|             | In lbs/acre                                      | 100 | 100                                                                  | 100                                                                  |
| Crabgrass   | In lbs/1000 sq ft                                | 2.3 | 2.3                                                                  | 2.3                                                                  |
| Goosegrass  |                                                  |     |                                                                      |                                                                      |

**CRABGRASS CONTROL**

For pre-emergent activity, **SPECTICLE G** is not effective until activated by rainfall or irrigation. **SPECTICLE G** applied to crabgrass that has emerged will not provide crabgrass control.

**GOOSEGRASS CONTROL**

**SPECTICLE G** provides preemergent control of goosegrass emerging from seed. For pre-emergent activity, **SPECTICLE G** is not effective until activated by rainfall or irrigation. **SPECTICLE G** does not control perennial (continual growth) goosegrass or plants emerging from existing crowns. If goosegrass is evident at time of application, use a labeled post emergent herbicide for control of existing plants.

**ANNUAL BLUEGRASS CONTROL**

Make pre-emergent applications of **SPECTICLE G** between August and November depending on the germination of annual bluegrass in each geographical area. The optimum timing is dependent on peak annual bluegrass germination and local conditions in a particular year.

**BROADLEAF WEED CONTROL**

**SPECTICLE G** provides preemergence control of many broadleaf weeds. Control is affected by the size of the seed, and the depth of the seed within the soil profile. **SPECTICLE G** must be activated by rainfall or irrigation prior to germination. Large seeded broadleaf weeds or weeds germinating deeper than several inches in the soil profile may not be effectively controlled by **SPECTICLE G**.

**ANNUAL SEDGES AND ANNUAL KYLLINGA CONTROL**

**SPECTICLE G** provides preemergence control of sedges and kyllinga emerging from seed. **SPECTICLE G** does not control established sedges and kyllinga (green kyllinga), or sedges emerging from tubers (nutlets) including yellow and purple nutedge.

**Weeds Controlled or Suppressed by SPECTICLE G\*\***

| Broadleaf Weeds             |                              |                           |                             |
|-----------------------------|------------------------------|---------------------------|-----------------------------|
| American black nightshade   | <i>Solanum americanum</i>    | Dandelion, common         | <i>Taraxacum officinale</i> |
| Bittercress                 | <i>Cardamine</i> sp.         | Eclipta                   | <i>Eclipta alba</i>         |
| Black medic*                | <i>Medicago lupulina</i>     | Evening primrose, Common  | <i>Oenothera biennis</i>    |
| Buckweed, wild              | <i>Polygonum convolvulus</i> | Evening primrose, Cutleaf | <i>Oenothera laciniata</i>  |
| California burclover        | <i>Medicago polymorpha</i>   | False chamomile*          | <i>Matricaria maritima</i>  |
| Canada thistle, Common      | <i>Cirsium arvense</i>       | Filaree, Redstem          | <i>Erodium cicutarium</i>   |
| Chickweed, Common           | <i>Stellaria media</i>       | Florida pusley            | <i>Richardia scabra</i>     |
| Chickweed, Mouse-ear        | <i>Cerastium vulgatum</i>    | Groundsel, Common         | <i>Senecio vulgaris</i>     |
| Clover, White               | <i>Trifolium repens</i>      | Hairy fleabane            | <i>Erigeron bonariensis</i> |
| Corn speedwell              | <i>Veronica arvensis</i>     | Hairy nightshade          | <i>Solanum sarrachoides</i> |
| Cudweed, Linear-leaf/purple | <i>Gnaphalium purpureum</i>  | Henbit                    | <i>Lamium amplexicaule</i>  |
| Curly dock                  | <i>Rumex crispus</i>         | Horseweed/ Marestail      | <i>Conyza canadensis</i>    |
| Dandelion, Cat's ear        | <i>Hypochoeris radicata</i>  | Kochia                    | <i>Kochia scoparia</i>      |

Weeds Controlled or Suppressed by SPECTICLE G\*\* (continued)

| Broadleaf Weeds               |                                 |                           |                                |
|-------------------------------|---------------------------------|---------------------------|--------------------------------|
| Lambsquarters, Common         | <i>Chenopodium album</i>        | Spurge, Spotted           | <i>Euphorbia maculata</i>      |
| Lawn burweed                  | <i>Soliva sessilis</i>          | Tassel flower             | <i>Emilia sonchifolia</i>      |
| Little mallow                 | <i>Malva parviflora</i>         | Redmaids                  | <i>Calandrinia caulescens</i>  |
| London rocket*                | <i>Sisymbrium irio</i>          | Sesbania, Hemp*           | <i>Sesbania exaltata</i>       |
| Longstalked phyllanthus       | <i>Phyllanthus tenellus</i>     | Shepherd's-purse          | <i>Capsella bursa-pastoris</i> |
| Mulberry weed                 | <i>Fatoua villosa</i>           | Sida, Prickly/Teaweed*    | <i>Sida spinosa</i>            |
| Mustard, Black*               | <i>Brassica nigra</i>           | Southern brassbuttons*    | <i>Cotula australis</i>        |
| Mustard, Wild                 | <i>Sinapis arvensis</i>         | Sowthistle, Annual        | <i>Conchus oleraceus</i>       |
| Parthenium                    | <i>Parthenium hysterophorus</i> | Sunflower*, Common        | <i>Helianthus annuus</i>       |
| Pigweed, Prostrate            | <i>Amaranthus blitoides</i>     | Swinecress                | <i>Coronopus didymus</i>       |
| Pigweed, Redroot              | <i>Amaranthus retroflexus</i>   | Tropic ageratum           | <i>Ageratum conyzoides</i>     |
| Plantain, Buckhorn            | <i>Plantago lanceolata</i>      | Velvetleaf*               | <i>Abutilon theophrasti</i>    |
| Plantain, Paleseed            | <i>Plantago virginica</i>       | Vetch, Purple*            | <i>Vicia benghalensis</i>      |
| Prostrate knotweed            | <i>Polygonum aviculare</i>      | Wild carrot*              | <i>Daucus carota</i>           |
| Prostrate spurge              | <i>Euphorbia maculata</i>       | Willowherb                | <i>Epilobium brachycarpum</i>  |
| Puncturevine                  | <i>Tribulus terrestris</i>      | Woodsorrel, Yellow*       | <i>Oxalis stricta</i>          |
| Purslane, Common              | <i>Portulaca oleracea</i>       | Woodsorrel/Oxalis*        | <i>Oxalis sp</i>               |
| Ragweed, Common*              | <i>Ambrosia artemisiifolia</i>  |                           |                                |
| GRASSES, MONOCOTS, AND SEDGES |                                 |                           |                                |
| Annual bluegrass/ 'Poa'       | <i>Poa annua</i>                | Goosegrass                | <i>Eleusine indica</i>         |
| Annual kylinga                | <i>Cyperus sesquiflorus</i>     | Guineagrass               | <i>Panicum maximum</i>         |
| Barneyardgrass, Common        | <i>Echinochloa crus-galli</i>   | Kyllinga, Fragrant/Annual | <i>Kyllinga odorata</i>        |
| Cheatgrass                    | <i>Bromus secalinus</i>         | Mouse barley              | <i>Hordeum murinum</i>         |
| Crabgrass                     | <i>Digitaria sp</i>             | Nutsedge, Purple*         | <i>Cyperus rotundus</i>        |
| Crabgrass, Henry              | <i>Digitaria ciliaris</i>       | Nutsedge, Yellow*         | <i>Cyperus esculentus</i>      |
| Crabgrass, Large/Hairy        | <i>Digitaria sanguinalis</i>    | Rice flatsedge            | <i>Cyperus iria</i>            |
| Crabgrass, Smooth             | <i>Digitaria ischaemum</i>      | Ryegrass, Italian         | <i>Lolium multiflorum</i>      |
| Doveweed                      | <i>Murdannia nudiflora</i>      | Ryegrass, Perennial       | <i>Lolium perenne</i>          |
| Foxtail brome                 | <i>Bromus rubens</i>            | Sandbur                   | <i>Cenchrus longispinus</i>    |
| Foxtail, Giant                | <i>Setaria faberi</i>           | Sedge, Annual***          | <i>Cyperus spp.</i>            |
| Foxtail, Green                | <i>Setaria viridis</i>          | Tufted lovegrass          | <i>Eragrostis pectinacea</i>   |
| Foxtail, Yellow               | <i>Pennisetum glaucum</i>       |                           |                                |

\* Weeds suppressed by **SPECTICLE G**  
\*\*Higher labeled rates should be used if weed pressure is historically high or longer residual control is desired (See Use Rates in the **SPECTICLE G** chart). The rate chosen should not exceed the tolerance for an individual turf type. Weed control can also be achieved with multiple applications of **SPECTICLE G**. Do not exceed a total of 400 lbs of product per acre per year (9.2 lbs/1000 sq ft/year) with multiple applications.  
\*\*\***SPECTICLE G** only controls sedges and kyllinga emerging from seed. It does not control perennial sedge emerging from nutlets or perennial forms of kyllinga.

SEEDING, OVERSEEDING, RE-SEEDING, SPRIGGING AND SODDING

**SEEDING, RE-SEEDING, AND OVERSEEDING**  
Timing of seeding, re-seeding, overseeding, sprigging, and sodding turf relative to an application of **SPECTICLE G** needs to be tracked carefully. **SPECTICLE G** can inhibit root development, as well as the emergence of seed. Newly emerged seedlings may be damaged and establishment of sod may be affected if **SPECTICLE G** is applied to turf that is not well established.  
**Application of SPECTICLE G prior to overseeding bermudagrass with perennial ryegrass or reseeded into warm season turf**

**SPECTICLE G** may be used to control weeds in bermudagrass prior to overseeding with a cool season grass or re-seeding seeded cultivars of bermudagrass, provided that the interval between application and seeding is appropriate (See **INTERVAL BETWEEN APPLICATION OF SPECTICLE G AND SEEDING OR OVERSEEDING** table below.) Overseeding and re-seeding intervals are affected by the rate of **SPECTICLE G** applied, rainfall in a given year and soil texture. Applications made sooner than the suggested intervals may decrease the establishment of the new seedlings and reduce turf coverage.

| INTERVAL BETWEEN APPLICATION OF SPECTICLE G AND SEEDING OR OVERSEEDING |                              |
|------------------------------------------------------------------------|------------------------------|
| RATE OF SPECTICLE G                                                    | MINIMUM OVERSEEDING INTERVAL |
| 100-125 lbs/acre<br>2.3-2.9 lbs/1000 sq ft                             | 10 months                    |
| 125-200 lbs/acre<br>2.9-4.6 lbs/1000 sq ft                             | 12 months                    |

## **WARM SEASON GRASSES ESTABLISHED WITH SPRIGS**

**SPECTICLE G** may be applied to sprigged warm season turf. The sprigs need to be well rooted prior to application. Apply the herbicide no sooner than 16 months after sprigging to allow for good stand establishment. Prior to application, check rooting to make sure new roots are developing. If roots are not growing, delay application of **SPECTICLE G**.

## **SODDING WARM SEASON GRASSES**

Delay sodding into bareground or turf treated with **SPECTICLE G** until 6 months after application. Sodding before 6 months may inhibit stand establishment and reduce turf quality. For maximum establishment of sod, follow the directions for the use of activated charcoal prior to laying sod.

Delay the application of **SPECTICLE G** if sod is not established. Sod needs to be established at least 3 months prior to an application of **SPECTICLE G**.

## **ORNAMENTALS**

**SPECTICLE G** may be applied for pre-emergent weed control in landscape ornamentals. Apply **SPECTICLE G** prior to weed seed germination to dormant or actively growing ornamentals as listed in the table below. Apply **SPECTICLE G** to established ornamentals. If uncertain about ornamental establishment after transplanting, wait 2 months before applying **SPECTICLE G**.

### **USE RESTRICTIONS FOR SPECTICLE G ON LANDSCAPE ORNAMENTALS**

- Do not exceed the maximum single application rates specified under each ornamental use.
- Do not exceed a total of 400 lbs of **SPECTICLE G** per acre (9.2 lbs/1000 sq ft) in a 12-month period.
- Do not allow granules applied "over-the-top" to remain in contact with foliage. **SPECTICLE G** may cause localized injury to the foliage, especially young leaf tissue. If the granules should be in contact with the foliage after an application, wash off affected foliage immediately.
- Do not use **SPECTICLE G** on ornamentals where granules may become trapped in developing leaves or in meristematic areas (e.g. whorls of grasses and perennials such as hosta).
- Do not use **SPECTICLE G** around bedding plants (including annuals and herbaceous plants not specified, or not listed as tolerant on this label).
- Do not use **SPECTICLE G** in landscape beds within 12 months prior to seeding or 3 months prior to planting.
- Do not use **SPECTICLE G** within the dripline of bearing fruit and nut trees. Non-bearing trees are defined as trees that will not bear fruit until at least 1 year after treatment.
- Do not use **SPECTICLE G** around non-bearing trees or vines unless they are at least 1 year old (citrus) and 3 years old (all others).
- Do not mix **SPECTICLE G** into the soil or potting mix. Mixing **SPECTICLE G** into the potting mix may inhibit developing roots.
- Do not cultivate or disturb the soil surface after application of **SPECTICLE G** as this may reduce weed control.
- If transplanting mature plants listed as tolerant on this label into soil treated with **SPECTICLE G** within the preceding 12 months, replace existing soil around roots to minimize effects on plant establishment.
- Do not apply **SPECTICLE G** to ornamental bulbs in landscapes or residential sites while they are emerging or after they have emerged.
- Do not apply **SPECTICLE G** to herbaceous perennials on this label while they are emerging or after they have emerged.
- Do not apply **SPECTICLE G** over-the-top of a deciduous ornamental at bud break.

### **LANDSCAPE ORNAMENTAL USES**

**SPECTICLE G** may be used in residential, commercial, state, and local plantings of ornamentals for preemergent weed control. **SPECTICLE G** should be applied only to established (rooted) plants and not to newly rooted cuttings/or seedlings. To avoid root damage, apply **SPECTICLE G** around transplants when the soil has firmly settled around the root area. Light irrigation or rainfall will help to settle the soil and seal surface cracks. For best results, remove existing mulch and replace mulch after an application of **SPECTICLE G**. If **SPECTICLE G** should contact foliage, brush or wash off to avoid damage. Tender annuals and herbaceous perennials may be sensitive to **SPECTICLE G**. Applications of **SPECTICLE G** should only be made to ornamentals listed on this label.

**AMOUNT TO USE:** Apply **SPECTICLE G** at 100-200 lbs of product per acre (2.3-4.6 lbs/1000 sq ft) around ornamentals.

To activate **SPECTICLE G** for maximum herbicidal benefit, irrigate the area treated with **SPECTICLE G** to move the herbicide into the soil within several days after application. With dry soil, use 0.25 inches of irrigation water, and with soil at or greater than field capacity, the amount of irrigation water should be reduced. In either case, do not create conditions that cause visible run-off of irrigation water. Adequate rainfall following an application will negate the need for irrigation.

A subsequent application of **SPECTICLE G** can be made within 90 days after the initial application to extend weed control provided that the total **SPECTICLE G** applied does not exceed 400 lbs per acre (9.2 lbs/1000 sq ft) in a 12 month period.

Application of **SPECTICLE G** to heavily mulched landscape beds may result in reduced weed control. **SPECTICLE G** may be applied to lightly mulched area after the mulch has settled. For best results remove existing mulch and replace mulch after an application of **SPECTICLE G**.

### **SYMPTOMS OF SPECTICLE G INJURY ON ORNAMENTALS**

**SPECTICLE G** may injure sensitive ornamentals by damaging roots or leaves. Plants damaged by root uptake will appear stunted, deformed, and may not recover. If granules of **SPECTICLE G** are allowed to remain in contact with treated plants, leaf symptoms may appear as lesions, discoloration, or curling. Symptoms may appear within 2-4 weeks after application. Newly formed leaves may be undamaged. It is essential that the user assess the severity of any symptoms on cultivars not listed on this label, before proceeding with large-scale applications of **SPECTICLE G**.

### **APPLICATION OF SPECTICLE G ON HERBACEOUS PLANTS**

**SPECTICLE G** may injure many annuals, perennials, and foliage plants. The product should not be applied to these plants unless they are listed on this label as tolerant.

## ORNAMENTALS AND THEIR CULTIVARS TOLERANT TO SPECTICLE G

Tolerant ornamentals and their cultivars are listed in the following table. If a cultivar to be treated is not listed on this label, treat several plants of the cultivar at the single maximum use rate (200 lbs per acre (4.6 lbs/1000 sq ft)) and evaluate 1-2 months later for possible injury and acceptable tolerance. If a plant type (designated by genus and species) is not listed on this label, do not apply **SPECTICLE G**. The user assumes responsibility for application on cultivars not listed in this table.

| Common Name                  | Scientific Name                   | Cultivar                     |
|------------------------------|-----------------------------------|------------------------------|
| Arborvitae                   | <i>Thuja</i> sp.                  | Green Giant                  |
| Azalea                       | <i>Rhododendron</i> sp.           | Girard's Rose                |
|                              |                                   | Fashion                      |
|                              |                                   | VF 14                        |
|                              |                                   | Golden Torch                 |
|                              |                                   | Judge Solomon                |
| Azalea, Encore               | <i>Rhododendron</i> sp.           | Autumn Debutante             |
| Beautybush                   | <i>Kolkwitzia amabilis</i>        |                              |
| Big bluestem                 | <i>Andropogon gerardii</i>        |                              |
| Bleeding Heart Vine          | <i>Clerodendrum thomsoniae</i>    | Bleeding Heart               |
| Bluebird                     | <i>Caryopteris x clandonensis</i> | Dark Knight                  |
| Bougainvillea                | <i>Bougainvillea</i>              | Barbara Karst                |
| Boxwood                      | <i>Buxus microphylla</i>          | Green Beauty                 |
|                              |                                   | Baby Gem                     |
| Butterfly bush               | <i>Buddleia</i>                   | Nanho Blue                   |
| Camellia, Japanese           | <i>Camellia japonica</i>          | Margaret Heathcliff Pink     |
| Camellia, Little leaf        | <i>Camellia sasanqua</i>          | Cleopatra Pink               |
| Cedar, Atlantic white        | <i>Chamaecyparis thyoides</i>     |                              |
| Cedar, Japanese              | <i>Cryptomeria japonica</i>       | Black Dragon                 |
|                              |                                   | Yoshino                      |
| Crape myrtle                 | <i>Lagerstroemia indica</i>       | Sarah's Favorite             |
|                              |                                   | Pocomoke Pink                |
| Cypress, false               | <i>Chamaecyparis</i> sp.          | Gold Mops                    |
| Daphne                       | <i>Daphne caucasica</i>           | Summer Ice                   |
| Florida pipestem             | <i>Leucothoe populifolia</i>      |                              |
| Forsythia                    | <i>Forsythia</i>                  | Lynwood Gold, Golden Bell    |
| Gardenia                     | <i>Gardenia radicans</i>          | Radicans                     |
|                              |                                   | Mystery                      |
| Gardenia                     | <i>Gardenia jasminoides</i>       | Frostproof                   |
|                              |                                   | Gold Dust                    |
| Gold-Dust Plant              | <i>Aucuba japonica</i>            | Gold Dust                    |
| Hardy kiwi                   | <i>Actinidia arguta</i>           | Anna                         |
| Hawaiian snowbush            | <i>Breynia disticha</i>           |                              |
| Hibiscus, Chinese            | <i>Hibiscus rosa-sinensis</i>     | San Diego Red, President Red |
| Holly, Chinese               | <i>Ilex cornuta</i>               | Needlepoint, Burdordii Nana  |
| Holly, Gallberry             | <i>Ilex glabra</i>                | Densa                        |
| Holly, Meservae              | <i>Ilex meserveae</i>             | Blue Princess                |
| Holly, Sky pencil            | <i>Ilex</i> sp.                   | Sky Pencil                   |
| Indian hawthorn              | <i>Raphiolepis indica</i>         | Pink Lady                    |
| Japanese mock-orange         | <i>Pittosporum tobira</i>         | Variegata                    |
| Jasmine, Asiatic/Yellow Star | <i>Trachelospermum asiaticum</i>  | Minima                       |
| Jasmine, Winter              | <i>Jasminum nudiflorum</i>        |                              |
| Juniper, Bar Harbour         | <i>Juniperus horizontalis</i>     | Bar Harbour                  |
| Juniper, Blue Pacific        | <i>Juniperus conferta</i>         | Blue Pacific                 |
| Juniper, Blue Rug            | <i>Juniperus horizontalis</i>     | Blue Rug                     |
| Lilac                        | <i>Syringa</i> x 'Penda'          | Bloomerang                   |
| Nandina                      | <i>Nandina</i>                    | Firepower                    |
| Oleander                     | <i>Nerium oleander</i>            | Cardinal Red                 |
| Pittosporum                  | <i>Pittosporum tobira</i>         | Variegata                    |
| Podocarpus                   | <i>Podocarpus macrophyllus</i>    |                              |
| Potentilla                   | <i>Potentilla fruticosa</i>       | Goldfinger                   |

Continued

| Common Name                     | Scientific Name                | Cultivar                     |
|---------------------------------|--------------------------------|------------------------------|
| Rose                            | <i>Rosa</i> sp.                | Pink Knockout®,<br>Knockout® |
| Rose, Virginia                  | <i>Rosa virginiana</i>         |                              |
| Rose mallow                     | <i>Hibiscus moscheutos</i>     |                              |
| Rose of Sharon                  | <i>Hibiscus syriacus</i>       | Pink Heart                   |
| Russian olive                   | <i>Elaeagnus angustifolia</i>  |                              |
| Slender deutzia                 | <i>Deutzia gracilis</i>        | Nikko                        |
| Sweet olive, fragrant tea olive | <i>Osmanthus fragrans</i>      |                              |
| Switchgrass                     | <i>Panicum virgatum</i>        | Shenandoah                   |
| Thin-fruit sedge                | <i>Carex flaccosperma</i>      |                              |
| Viburnum, Burkwood              | <i>Viburnum x burkwoodii</i>   |                              |
| Viburnum, Popcorn               | <i>Viburnum plicatum</i>       | Popcorn                      |
| Virginia sweetspire             | <i>Itea virginica</i>          |                              |
| Wax myrtle, Southern            | <i>Myrica cerifera</i>         |                              |
| Witchalder                      | <i>Fothergilla gardenii</i>    |                              |
| Yew podocarpus                  | <i>Podocarpus macrophyllus</i> |                              |

**Do not use SPECTICLE G on any of these plants or their cultivars; injury will occur.**

| Common Name            | Scientific Name                   |
|------------------------|-----------------------------------|
| Astilbe                | <i>Astilbe chinensis</i>          |
| Barberry               | <i>Berberis thunbergii</i>        |
| Blue fescue grass      | <i>Festuca glauca</i>             |
| Columbine              | <i>Aquilegia canadensis</i>       |
| Coneflower             | <i>Echinacea purpurea</i>         |
| Croton                 | <i>Codiaeum variegatum</i>        |
| Dewdrops               | <i>Duranta erecta</i>             |
| Dianthus               | <i>Dianthus</i> sp.               |
| Eulelia grass          | <i>Miscanthus sinensis</i>        |
| Euonymus, Japanese     | <i>Euonymus japonicus</i>         |
| Feather reed grass     | <i>Calamagrostis x acutiflora</i> |
| Fountain grass         | <i>Pennisetum alopecuroides</i>   |
| Fountain grass, Purple | <i>Pennisetum setaceum</i>        |
| Hydrangea              | <i>Hydrangea macrophylla</i>      |
| Ixora                  | <i>Ixora coccinea</i>             |

## NON-SELECTIVE USES

### HARDSCAPES

**SPECTICLE G** may be used to maintain bare ground in non-crop areas. These include gravel areas, rocky/sandy areas found in desert climates and non-paved paths on golf courses and in residential areas. For optimum weed control, remove all plant debris from the area to be treated. Irrigation or rainfall after application will provide maximum weed control. Apply 400 lbs of **SPECTICLE G** per acre (9.2 lbs/1000 sq ft) for these uses.

Observe all restrictions and tolerances for applications of **SPECTICLE G** around ornamentals listed on this label.

## IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury or ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSOURCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSOURCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR

BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer (reg'd), the Bayer Cross (reg'd), Specticle™ and Backed by Bayer™ are trademarks of Bayer. Knock Out® a registered trademark of CP (Delaware) Inc.



Produced for:

**Bayer Environmental Science**

A Division of Bayer CropScience LP

2 T. W. Alexander Drive

Research Triangle Park, NC 27709